

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074042 A2

(51) International Patent Classification: Not classified

[US/US]; 46 Lakeview Terrace, Watchung, NJ 07069 (US). **TIANLING CHEN, TianLing** [US/US]; 38 Park Street, #18C, Florham Park, NJ 07932 (US).

(21) International Application Number: PCT/EP02/02977

(22) International Filing Date: 18 March 2002 (18.03.2002)

(74) Agent: **BECKER, Konrad**; Novartis AG, Corporate Intellectual Property, Patent & Trademark Department, CH-4002 Basel (CH).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/277,153 19 March 2001 (19.03.2001) US
60/277,207 20 March 2001 (20.03.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NO, NZ, OM, PH, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TN, TR, TT, UA, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except AT, US*): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(71) Applicant (*for AT only*): **NOVARTIS-ERFINDUNGEN VERWALTUNGSGESELLSCHAFT M.B.H.** [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

Published:

— *without international search report and to be republished upon receipt of that report*

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ROTHERMEL, John, David** [US/US]; 46 School House Road, Randolph, NJ 07869 (US). **SCHRAN, Horst, F.** [US/US]; 17 Junard Drive, Morristown, NJ 07960 (US). **GREELEY, Diane**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/074042 A2

(54) Title: COMBINATIONS COMPRISING AN ANTIDIARRHEAL AGENT AND AN EPOTHILONE OR AN EPOTHILONE DERIVATIVE


(57) Abstract: Epothilone derivatives are co-administered with an antidiarrheal agent, e.g., a DPP-IV inhibitor, in the treatment of a proliferative disease.

Combinations Comprising an Antidiarrheal Agent and an Epothilone or an Epothilone Derivative

The invention relates to a pharmaceutical combination which comprises (a) an antidiarrheal agent, in particular, a dipeptidylpeptidase – IV (DPP-IV) inhibitor, (b) an epothilone derivative of formula I, and optionally at least one pharmaceutically acceptable carrier for simultaneous, separate or sequential use, in particular for the treatment of a proliferative disease, especially a solid tumor disease; a pharmaceutical composition comprising such a combination; the use of such a combination for the preparation of a medicament for the treatment of a proliferative disease; a commercial package or product comprising such a combination as a combined preparation for simultaneous, separate or sequential use; and to a method of treatment of a warm-blooded animal, especially a human.

The microtubule-stabilizing effect of epothilones was first described by Bollag et al., Cancer Research 55, 1995, 2325-33. A suitable treatment schedule of different types of tumors, especially tumors which are refractory to the treatment by other chemotherapeutics, in particular TAXOL™, is described in WO 99/43320.

The present invention pertains to a combination, such as a combined preparation or a pharmaceutical composition, which comprises (a) an antidiarrheal agent and (b) an epothilone derivative of formula I



(I)

in which compound A represents O or NR_N, wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, in which the active ingredients (a) and (b) are present in each case in free form or in the form of a pharmaceutically acceptable salt and

optionally at least one pharmaceutically acceptable carrier; for simultaneous, separate or sequential use.

A compound of formula I wherein A represents O, R is hydrogen and Z is O is known as epothilone A; a compound of formula I wherein A represents O, R is methyl and Z is O is known as epothilone B; a compound of formula I wherein A represents O, R is hydrogen and Z is a bond is known as epothilone C; a compound of formula I wherein A represents O, R is methyl and Z is a bond is known as epothilone D.

The term "a combined preparation", as used herein defines especially a "kit of parts" in the sense that the combination partners (a) and (b) as defined above can be dosed independently or by use of different fixed combinations with distinguished amounts of the combination partners (a) and (b), i.e., simultaneously or at different time points. The parts of the kit of parts can then, e.g., be administered simultaneously or chronologically staggered, that is at different time points and with equal or different time intervals for any part of the kit of parts. The ratio of the total amounts of the combination partner (a) to the combination partner (b) to be administered in the combined preparation can be varied, e.g. in order to cope with the needs of a patient sub-population to be treated or the needs of the single patient based on the severity of the diarrhea that the patient experiences.

The present invention especially relates to a combined preparation, which comprises (a) one or more unit dosage forms of an antidiarrheal agent and (b) one or more unit dosage forms of an epothilone derivative of formula I, especially epothilone B.

The antidiarrheal agent is administered to prevent, control or eliminate diarrhea that is sometimes associated with the administration of epothilones, especially epothilone B. Thus, the present invention also relates to a method of preventing or controlling diarrhea associated with administering an epothilone derivative of formula I, which comprises administering an effective amount of an antidiarrheal agent to the patient receiving treatment with the epothilone derivative.

The term "solid tumor" especially means breast cancer, ovarian cancer, cancer of the colon and generally the GI tract, cervix cancer, lung cancer, in particular small-cell lung cancer, and non-small-cell lung cancer, head and neck cancer, bladder cancer, cancer of the

prostate or Kaposi's sarcoma. The present combination inhibits the growth of solid tumors, but also liquid tumors. Furthermore, depending on the tumor type and the particular combination used a decrease of the tumor volume can be obtained. The combinations disclosed herein are also suitable to prevent the metastatic spread of tumors and the growth or development of micrometastases.

The structure of the active agents identified by code nos., generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g. Patents International (e.g. IMS World Publications). The corresponding content thereof is hereby incorporated by reference.

It will be understood that references to the combination partners (a) and (b) are meant to also include the pharmaceutically acceptable salts. If these combination partners (a) and (b) have, for example, at least one basic center, they can form acid addition salts. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The combination partners (a) and (b) having an acid group (for example COOH) can also form salts with bases. The combination partner (a) or (b) or a pharmaceutically acceptable salt thereof may also be used in form of a hydrate or include other solvents used for crystallization.

Epothilone derivatives of formula I wherein A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl and Z is O or a bond, and methods for the preparation of such epothilone derivatives are in particular generically and specifically disclosed in the patents and patent applications WO 93/10121, US 6,194,181, WO 98/25929, WO 98/08849, WO 99/43653, WO 98/22461 and WO 00/31247 in each case in particular in the compound claims and the final products of the working examples, the subject-matter of the final products, the pharmaceutical preparations and the claims is hereby incorporated into the present application by reference to this publications. Comprised are likewise the corresponding stereoisomers as well as the corresponding crystal modifications, e.g. solvates and polymorphs, which are disclosed therein.

The transformation of epothilone B to the corresponding lactam is disclosed in Scheme 21 (page 31, 32) and Example 3 of WO 99/02514 (pages 48 - 50). The transformation of a compound of formula I which is different from epothilone B into the corresponding lactam

can be accomplished analogously. Corresponding epothilone derivatives of formula I wherein R_N is lower alkyl can be prepared by methods known in the art such as a reductive alkylation reaction starting from the epothilone derivative wherein R_N is hydrogen.

Epothilone derivatives of formula I, especially epothilone B, can be administered as part of pharmaceutical compositions which are disclosed in WO 99/39694.

In a specific embodiment, the epothilone derivative is a compound of formula I in which compound A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond.

In the epothilone derivative of formula I preferably A represents O, R is lower alkyl, e.g. ethyl or, most preferably, methyl and Z is preferably O.

Antidiarrheal agents and protocols for their administration are known to those skilled in the art. Antidiarrheal agents suitable for use in the inventive methods and compositions include, but are not limited to, natural opioids, such as tincture of opium, paregoric, and codeine, synthetic opioids, such as diphenoxylate, difenoxin and loperamide, bismuth subsalicylate, lanreotide, vapreotide and octreotide, motilin antagonists, COX2 inhibitors like celecoxib, glutamine, thalidomide and traditional antidiarrheal remedies, such as kaolin, pectin, berberine and muscarinic agents. In one embodiment of the invention, the antidiarrheal agent is selected from codeine, tincture of opium, paregoric, diphenoxylate, difenoxine and loperamide. In another embodiment of the invention, the antidiarrheal agent is selected from lanreotide, vapreotide and octreotide. Of these three compounds, octreotide is especially preferred. The latter compound or its acetate can be obtained and employed as described in US 4,395,403, or, e.g., in the form of its acetate or pamoate as described in US 5,538,739. In particular, octreotide can be administered to the patient as marketed under the trademarks SANDOSTATINTM and SANDOSTATIN LARTM.

The antidiarrheal agent employed in the present invention can also be a DPP-IV inhibitor. DPP-IV inhibitors are known in the art for the treatment of diabetes. Protocols for their administration in the treatment of diabetes are known to those skilled in the art. However, in the present invention the DPP-IV inhibitor functions to prevent and/or control diarrhea that is sometimes associated with the administration of epothilone derivatives.

DPP-IV is responsible for inactivating GLP-1. More particularly, DPP-IV generates a GLP-1 receptor antagonist and thereby shortens the physiological response to GLP-1. GLP-1 is a major stimulator of pancreatic insulin secretion and has direct beneficial effects on glucose disposal.

The DPP-IV inhibitor can be peptidic or non-peptidic. Preferably, the DPP-IV inhibitor is non-peptidic.

Preferably, DPP-IV inhibitors are employed in the present invention which are generically and specifically disclosed in WO 98/19998, DE 196 16 486 A1, WO 00/34241 and WO 95/15309, in each case in particular in the compound claims and the final products of the working examples, the subject-matter of the final products, the pharmaceutical preparations and the claims are hereby incorporated into the present application by reference to these publications. DPP728 and LAF237 are specifically disclosed in Example 3 of WO 98/19998 and Example 1 of WO 00/34241, respectively. A further suitable DPP-IV inhibitor of formula VI (see below) is specifically described in Diabetes 1998, 47, 1253-1258. DPP728 can be formulated as described on page 20 of WO 98/19998.

N-Peptidyl-O-aryl hydroxylamines, e.g. of formula VII or VIIa (see below), and their preparation are described by H.U. Demuth et al. in J. Enzyme Inhibition 1988, Vol. 2, pages 129-142, especially on pages 130-132.

Unless stated otherwise in the present disclosure organic radicals designated "lower" contain not more than 7, preferably not more than 4, carbon atoms and the following expressions have the meanings as given below:

Halogen represents preferably fluoro, chloro or bromo.

Lower alkyl is, if not stated otherwise, preferably ethyl or, most preferably, methyl. (C₁₋₈)Alkyl is branched or preferably unbranched alkyl, preferably lower alkyl, e.g. methyl or ethyl.

Lower alkylene is preferably methylene, ethylene or propylene. It can be unsubstituted or substituted e.g. by hydroxy.

Lower alkoxy is preferably methoxy or ethoxy. (C₂₋₄)Alkoxy is e.g. ethoxy or propoxy.

Cycloalkyl is e.g. C₃-C₁₂cycloalkyl, preferably cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclodecyl; or bicycloalkyl such as bicycloheptyl. Cycloalkenyl is preferably 1-cyclohexenyl, 2-cyclohexenyl, 3-cyclohexenyl, 1-cyclopentenyl or 1-cyclopentenyl.

(C₁₋₃)Hydroxyalkyl is e.g. 3-hydroxypropyl, 1-hydroxyethyl or hydroxymethyl.


C₄-C₆-Alkylenimino which is unsubstituted or substituted by one or two lower alkyl groups is, for example, pyrrolidinyl, methylpyrrolidinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 2-methyl-1-piperidinyl or hexamethylenimino. Preferably, C₄-C₆-alkylenimino is 1-piperidinyl.

A [3.1.1]bicyclic carbocyclic moiety optionally substituted as defined above preferably is bicyclo[3.1.1]hept-2-yl optionally disubstituted in 6-position with methyl, or bicyclo[3.1.1]-hept-3-yl optionally trisubstituted with one methyl in 2-position and two methyl groups in 6-position. A [2.2.1]bicyclic carbocyclic moiety optionally substituted as defined above preferably is bicyclo[2.2.1]hept-2-yl.

Aryl comprises preferably 6 to 12 carbon atoms and is e.g. phenyl, tolyl or naphthyl, each of which can be substituted e.g. by lower alkyl or halogen.

The term "heteroaryl" refers to an aromatic heterocyclic radical selected, for example, from the group consisting of pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazoliny, imidazolidinyl, oxazolyl, oxazolidinyl, isoxazoliny, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, azepiny, 4-piperidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranyl, morpholiny, thiamorpholiny, thiamorpholiny sulfoxide, thiamorpholiny sulfone, 1,3-dioxolane, indolyl, benzothiazolyl, benzoxazolyl, benzothieryl, quinuclidinyl, quinoliny, tetrahydroisoquinoliny, isoquinoliny, benzimidazolyl, benzopyranyl, indoliziny, benzofuryl, chromonyl, coumariny, benzopyranyl, cinnoliny, quinoxaliny, indazolyl, pyrrolopyridyl, furopyridiny, dihydrobenzothiazolyl, dihydroisoindolyl, dihydroquinazoliny and tetrahydroquinazoliny.

Preferred DPP-IV inhibitors are N-(N'-substituted glycyI)-2-cyanopyrrolidines represented by formula (I),



wherein R is:

a) $R_1R_{1a}N(CH_2)_m$ - wherein

R_1 is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with lower alkyl, lower alkoxy, halogen, trifluoromethyl, cyano or nitro; or phenyl optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen;

R_{1a} is hydrogen or (C_{1-8}) alkyl; and

m is 2 or 3;

b) (C_{3-12}) cycloalkyl optionally monosubstituted in the 1-position with (C_{1-3}) hydroxyalkyl;

c) $R_2(CH_2)_n$ - wherein either

R_2 is phenyl optionally mono- or independently di- or independently trisubstituted with lower alkyl, lower alkoxy, halogen or phenylthio optionally monosubstituted in the phenyl ring with hydroxymethyl; or is (C_{1-8}) alkyl; a [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted with (C_{1-8}) alkyl; a pyridinyl or naphthyl moiety optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen; cyclohexene; or adamantyl; and

n is 1 to 3; or

R_2 is phenoxy optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen; and

n is 2 or 3;

d) $(R_3)_2CH(CH_2)_2$ - wherein each R_3 independently is phenyl optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen;

e) $R_4(CH_2)_p$ - wherein R_4 is 2-oxopyrrolidinyl or (C_{2-4}) alkoxy and p is 2 to 4;

f) **isopropyl** optionally monosubstituted in 1-position with (C_{1-3}) hydroxyalkyl;

g) R_5 wherein R_5 is: indanyl; a pyrrolidinyl or piperidinyl moiety optionally substituted with benzyl; a [2.2.1]- or [3.1.1]bicyclic carbocyclic moiety optionally mono- or plurisubstituted

with (C₁₋₈)alkyl; adamantyl; or (C₁₋₈)alkyl optionally mono- or independently plurisubstituted with hydroxy, hydroxymethyl or phenyl optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen;

h) a substituted adamantyl

in free form or in acid addition salt form.

In a preferred embodiment of the invention, the N-(N'-substituted glycyI)-2-cyanopyrrolidine is represented by formula (I), wherein

R is **R₁R_{1a}N(CH₂)_m**- wherein

R₁ is a pyridinyl or pyrimidinyl moiety optionally mono- or independently disubstituted with lower alkyl, lower alkoxy, halogen, trifluoromethyl, cyano or nitro; or phenyl optionally mono- or independently disubstituted with lower alkyl, lower alkoxy or halogen;

R_{1a} is hydrogen or (C₁₋₈)alkyl; and

m is 2 or 3;

in free form or in acid addition salt form.

More preferably, the N-(N'-substituted glycyI)-2-cyanopyrrolidine is represented by formula (I), wherein

R is **R₁R_{1a}N(CH₂)_m**- wherein

R₁ is a pyridinyl moiety optionally mono- or independently disubstituted with lower alkyl, lower alkoxy, halogen, trifluoromethyl, cyano or nitro;

R_{1a} is hydrogen or (C₁₋₈)alkyl; and

m is 2 or 3;


in free form or in acid addition salt form.

Most preferably, the N-(N'-substituted glycyI)-2-cyanopyrrolidine of formula I is (S)-1-{2-[5-cyanopyridin-2-yl]amino}ethyl-aminoacetyl-2-cyano-pyrrolidine (DPP728) or (S)-1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-pyrrolidine (LAF237).


In another preferred embodiment, the DPP-IV inhibitor is selected from the compounds of formulae II, III, IV and V:

A - B


(formula II, groups G1 and G2)



(formula III, group G3)




(formula IV, group G3) and



(formula V, group G3)

where B is



f is 1 or 2;

g is 0, 1 or 2;


X is CH₂, O, S, SO, SO₂, NH or NR_{α1}, where R_{α1} is lower alkyl (C₁ to C₆);

-Y is -N, -CH or -C= (when the -CO group of A is replaced with -CH= or -CF=);

R_α is H, CN, CHO, B(OH)₂, PO₃H or an ester thereof, CC-R_{α7}, or CH=N-R_{α8} where R_{α7} is H, F, lower alkyl (C₁ to C₆), CN, NO₂, OR_{α9}, CO₂R_{α9} or COR_{α9}; R_{α9} is lower alkyl (C₁ to C₆); R_{α8} is Ph, OH, OR_{α9}, OCOR_{α9} or OBn; A is attached to Y;

and wherein for the group G1 compounds

(a) when R_α is H, A is an α-amino-acyl group derived from an α-amino-acid bearing a cycloaliphatic side-chain or is a β-amino-acyl group of general formula




where h is 1 to 6, the ring in either case optionally having unsaturation and/or heteroatom substitution;


(b) when $R\alpha$ is CN, $CC-R\alpha_7$, or $CH=N-R\alpha_8$, A is as defined at (a) and in addition may be derived from any L- α -amino acid bearing a lipophilic side-chain;

(c) and when $R\alpha$ is CHO or $B(OH)_2$, A is a β -amino-acyl group as defined under (a);


for the group G2 compounds, $R\alpha$ is H, CN, $C=C-R\alpha_7$ or $-CH=N-R\alpha_8$ and A is



where a is 1 - 5; D^1 is $-G-(CH_2)_b-(R\alpha_4)_q-R\alpha_3$; G is O, NH or NMe; b is 0 - 12; q is 0 - 5; D^2 is D^1 with $G \neq O$; $R\alpha_4$ is $Z-NH-(CH_2)_c-$ or $NH-Z-(CH_2)_c-$ where c is 1 - 12 and Z is CO, CH_2 or SO_2 ; $R\alpha_3$ is CO_2H or an ester thereof, $CONH_2$, $CONHNH_2$, $CONR\alpha_5R\alpha_6$, $CONHNR\alpha_5R\alpha_6$, PO_3H or an ester thereof, SO_3H , SO_2NH_2 , $SO_2NR\alpha_5R\alpha_6$, OH, $OR\alpha_5$, substituted or unsubstituted aryl or heteroaryl, NH_2 , $NR\alpha_5R\alpha_6$, $NHCO_2R\alpha_5$, $NHSO_2NR\alpha_5R\alpha_6$, $NHCOR\alpha_5$, $NH-SO_2R\alpha_5$, $NH-CH(:NR\alpha_5)NR\alpha_5R\alpha_6$, $NHCONHR\alpha_5R\alpha_6$, sugar, CO-aminosugar, $NHCO$ -aminosugar or $-NHCS$ -aminosugar; and $R\alpha_5$ and $R\alpha_6$ are independently selected from H and lower alkyl, fluoroalkyl and cycloalkyl group of up to 8 atoms and aryl, heteroaryl and alkyl heteroaryl groups of up to 11 atoms or $R\alpha_5$ and $R\alpha_6$ may together comprise a chain (C_3 to C_8); or is



where $R\alpha_{10}$ is H or Me, the ring may contain more heteroatoms, E is $J-(CH_2)_b-(R\alpha_4)_q-R\alpha_3$, J = CO, CH_2 or SO_2 , and a, b, q, $R\alpha_3$ and $R\alpha_4$ are as defined under (i); or is




where $R\alpha_2$ is H or Me, the ring may contain one or more heteroatoms, and L is $(CH_2)_d-(CO)_r$, $(CH_2)_b-(R\alpha_4)_q-R\alpha_3$ or $(CH_2)_e-NR\alpha_{10}-(CH_2)_b-(R\alpha_4)_q-R\alpha_3$ where r is 0 or 1, d is 0 - 4, e is 2 - 4, and b, q, $R\alpha_3$ and $R\alpha_4$ are as defined under (i);

and for the group G3 compounds, each B may have any identity defined therefor above, each A may be chosen from any group G2 structure (i), (ii) or (iii) above with the terminal groups $R\alpha_3$ in the A residues replaced with a shared group $-\epsilon-\omega-\epsilon-$ or $-\epsilon-\epsilon-$ or $-\omega-$, and ϵ and ω are selected independently from CH_2 , O, NH, CO, S, SO_2 , Ph and NHMe;

and wherein in groups G2 and G3 at least one CH_2 group in a chain may be replaced by a bioisostere thereof or any amide group which connects A and B in a group G1, G2 or G3 compound or which is in a side-chain of A in a Group G2 or G3 compound may be replaced by an amide bioisostere,

in free form or in acid addition salt form.

In another preferred embodiment, the DPP-IV inhibitor is a compound of formula VI




in free form or in acid addition salt form.

In a further preferred embodiment, the DPP-IV inhibitor is a N-peptidyl-O-aryl hydroxylamine or a pharmaceutically acceptable salt thereof. Aryl is, for example, naphthylcarbonyl; or benzoyl which is unsubstituted or mono- or disubstituted, for example, by lower alkoxy, lower alkyl, halogen or, preferably, nitro. The peptidyl moiety comprises preferably two α -

amino acids, e.g. glycine, alanine, leucine, phenylalanine, lysine or proline, of which the one attached directly to the hydroxylamine nitrogen atom is preferably proline.

Preferably, the N-peptidyl-O-aryl hydroxylamine is a compound of formula VII




wherein

j is 0, 1 or 2;

R_{E1} represents the side chain of a natural amino acid; and

R_{E2} represents lower alkoxy, lower alkyl, halogen or nitro;
or a pharmaceutically acceptable salt thereof.

In a very preferred embodiment of the invention, the N-peptidyl-O-aryl hydroxylamine is a compound of formula VIIa



or a pharmaceutically acceptable salt thereof.

In a highly preferred embodiment, the component (a) DPP-IV inhibitor is (S)-1-{2-[5-cyano-pyridin-2-yl)amino]ethyl-aminoacetyl}-2-cyano-pyrrolidine (DPP728), and the component (b) epothilone derivative is epothilone B.

In another highly preferred embodiment, the component (a) DPP-IV inhibitor is (S)-1-[(3-hydroxy-1-adamantyl)amino]acetyl-2-cyano-pyrrolidine (LAF237) and the component (b) epothilone derivative is epothilone B.

The present invention especially relates to a combined preparation, which comprises (a) one or more unit dosage forms of an DPP-IV inhibitor and (b) one or more unit dosage forms of an epothilone derivative of formula I, especially epothilone B.

A combination which comprises (a) an antidiarrheal agent and (b) an epothilone derivative of formula I in which compound A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, in which the active ingredients are present in each case in free form or in the form of a pharmaceutically acceptable salt and optionally at least one pharmaceutically acceptable carrier, will be referred to hereinafter as a COMBINATION OF THE INVENTION.

When the combination partners employed in the COMBINATION OF THE INVENTION are applied in the form as marketed as single drugs, their dosage and mode of administration can take place in accordance with the information provided on the package insert of the respective marketed drug in order to result in the beneficial effect described herein, if not mentioned herein otherwise.

The antidiarrheal agent is administered as a preventative measure throughout the cycle or as needed when diarrhea occurs.

In a preferred embodiment of the present invention, the subject receives the epothilone derivative of formula I once weekly for several weeks, for example three weeks, followed by one or several weeks off and the antidiarrheal agent is administered as a preventative measure by pretreating the subject with the antidiarrheal agent before the administration of the epothilone derivative begins and continuing administration of the antidiarrheal agent throughout the cycles, or by administering the antidiarrheal agent throughout the cycles without pretreatment or by administering antidiarrheal agent as needed when diarrhea occurs during the cycles, with or without a pretreatment. As an example, when the epothilone derivative is administered once weekly for three weeks with one week off, each four week interval will be considered one cycle.

An effective amount of the antidiarrheal agent is an amount sufficient to prevent, control, or eliminate diarrhea associated with the administration of an epothilone derivative, especially it

is an amount which increases the amount of the epothilone derivative that can administered when the diarrhea is the does limiting toxicity of the epothilone derivative, especially epothilone B.

The COMBINATION OF THE INVENTION can be a combined preparation or a pharmaceutical composition.

It is one objective of this invention to provide a pharmaceutical composition comprising a quantity, which is therapeutically effective against a proliferative disease comprising the COMBINATION OF THE INVENTION, i.e., in such pharmaceutical composition the combination partners (a) and (b) are administered together in a fixed combination.

The pharmaceutical compositions according to the invention can be prepared in a manner known per se and are those suitable for enteral, such as oral or rectal, and parenteral administration to mammals (warm-blooded animals), including man.

The novel pharmaceutical composition contain, for example, from about 10 % to about 100 %, preferably from about 20 % to about 60 %, of the active ingredients. Pharmaceutical preparations for the combination therapy for enteral or parenteral administration are, for example, those in unit dosage forms, such as sugar-coated tablets, tablets, capsules or suppositories, and furthermore ampoules. If not indicated otherwise, these are prepared in a manner known per se, for example by means of conventional mixing, granulating, sugar-coating, dissolving or lyophilizing processes. It will be appreciated that the unit content of a combination partner contained in an individual dose of each dosage form need not in itself constitute an effective amount since the necessary effective amount can be reached by administration of a plurality of dosage units.

In preparing the compositions for oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents; or carriers such as starches, sugars, microcrystalline cellulose, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations such as, for example, powders, capsules and tablets, with the solid oral preparations being preferred over the liquid preparations. Because of their

ease of administration, tablets and capsules represent the most advantageous oral dosage unit form in which case solid pharmaceutical carriers are obviously employed.

In particular, a therapeutically effective amount of each of the combination partner of the COMBINATION OF THE INVENTION may be administered separately, i.e., the components may be administered simultaneously or sequentially and in any order. For example, the method of treatment of a proliferative disease according to the invention may comprise (i) administration of the first combination partner in free or pharmaceutically acceptable salt form and (ii) administration of the second combination partner in free or pharmaceutically acceptable salt form, simultaneously or sequentially in any order, in jointly therapeutically effective amounts, preferably in synergistically effective amounts, e.g. in daily dosages corresponding to the amounts described herein. The individual combination partners of the COMBINATION OF THE INVENTION can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. One agents can, e.g., be an enteral formulation and the other can be administered parenterally. Furthermore, the term administering also encompasses the use of a pro-drug of a combination partner that convert *in vivo* to the combination partner as such. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment and the term "administering" is to be interpreted accordingly.

The effective dosage of each of the combination partners employed in the COMBINATION OF THE INVENTION may vary depending on the particular compound or pharmaceutical composition employed, the mode of administration, the condition being treated, the severity of the condition being treated. Thus, the dosage regimen the COMBINATION OF THE INVENTION is selected in accordance with a variety of factors including the route of administration and the renal and hepatic function of the patient. A physician, clinician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the single active ingredients required to prevent, counter or arrest the progress of the condition. Optimal precision in achieving concentration of the epothilone derivative within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the active ingredients' availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of the active ingredients.

If the the warm-blooded animal is a human, the dosage of a compound of formula I is preferably in the range of about 0.25 to 75, preferably 0.5 to 50, e.g. 2.5, mg/m² once weekly for two to four, e.g. three, weeks, followed by 6 to 8 days off in the case of an adult patient.

The antidiarrheal agent is preferably administered from one to four times per day according to established protocols for the antidiarrheal agent.

A DPP-IV inhibitor, if employed, is preferably administered according to known protocols for the treatment of diabetes. Preferably, the dose is in the range from 25 mg to 1000 mg per day.

Moreover, the present invention relates to a method of treating a warm-blooded animal having a proliferative disease comprising administering to the animal a COMBINATION OF THE INVENTION in a quantity which is therapeutically effective against a proliferative disease and which reduces any diarrhea associated with the administration of the epothilone derivative.

Furthermore, the present invention pertains to the use of a COMBINATION OF THE INVENTION for the treatment of a proliferative disease and for the preparation of a medicament for the treatment of a proliferative disease.

Moreover, the present invention provides a commercial package comprising as active ingredients COMBINATION OF THE INVENTION, together with instructions for simultaneous, separate or sequential use thereof in the treatment of a proliferative disease.

The following Examples illustrate the invention described above; they are not, however, intended to limit the scope of the invention in any way. The beneficial effects of the COMBINATION OF THE INVENTION can also be determined by other test models known as such to the person skilled in the pertinent art.

Example 1: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of a once a week administration of 2.5 mg epothilone B as a 5 minute bolus or as a 15 minute bolus for three weeks followed by 14

days of rest. Throughout the treatment the patient receives from 2 mg to 16 mg of loperamide daily as its hydrochloride salt to control diarrhea.

Example 2: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of 2.5 mg epothilone B administered as a 5 minute bolus once a week for three weeks followed by 14 days of rest. The patient receives up to 16 mg of loperamide as its hydrochloride salt to daily when diarrhea occurs.

Example 3: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of 2.5 mg epothilone B administered as a 5 minute bolus once a week for three weeks followed by 14 days of rest. Throughout the treatment the patient receives from one to six 50 mg doses of DPP728 daily to control diarrhea.


Example 4: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of 2.5 mg epothilone B administered as a 5 minute bolus once a week for three weeks followed by 14 days of rest. The patient receives up to 300 mg of DPP728 daily when diarrhea occurs.

Example 5: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of 2.5 mg epothilone B administered as a 15 minute bolus once a week for three weeks followed by 14 days of rest. Throughout the treatment the patient receives from one to six 50 mg doses of LAF 237 daily to control diarrhea.

Example 6: A human patient suffering from a proliferative disease is treated with 6 cycles of epothilone B wherein each cycle consists of 2.5 mg epothilone B administered as a 5 minute bolus once a week for three weeks followed by 14 days of rest. The patient receives up to 300 mg of LAF 237 daily when diarrhea occurs.


What is claimed is:

1. A combination which comprises (a) an antidiarrheal agent and (b) an epothilone derivative of formula I



wherein A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond,
in which the active ingredients (a) and (b) are present in each case in free form or in the form of a pharmaceutically acceptable salt and optionally at least one pharmaceutically acceptable carrier; for simultaneous, separate or sequential use.


2. Combination according to claim 1 comprising an epothilone derivative of formula I wherein A represents O, R is lower alkyl or hydrogen and Z is O or a bond.
3. Combination according to claim 1 or 2, wherein the antidiarrheal agent is a DPP-IV inhibitor.
4. Combination according to claim 1 or 2, wherein the antidiarrheal agent is a natural opioid, a synthetic opioid, bismuth subsalicylate, lanreotide, vapreotide, octreotide, a COX2 inhibitor, a motilin antagonist, kaolin, glutamine, thalidomide, pectin or berberine, or a pharmaceutically acceptable salt thereof.
5. A method of preventing or controlling diarrhea associated with administering an epothilone derivative of formula I



(I)

in which compound A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, in free form or in the form of a pharmaceutically acceptable salt, to a patient, which comprises administering an effective amount of an antidiarrheal agent to the patient receiving treatment with the epothilone derivative.


6. The method of claim 5 wherein the epothilone derivative is epothilone B.
7. A pharmaceutical composition comprising a quantity which is jointly therapeutically effective against a proliferative disease of a combination according to any one of claims 1 to 3 and at least one pharmaceutically acceptable carrier.
8. Use of a combination according to any one of claims 1 to 4 for the treatment of a proliferative disease.
9. Use of a combination according to any one of claims 1 to 4 for the preparation of a medicament for the treatment of a proliferative disease.
10. Use of an antidiarrheal agent in combination with an epothilone derivative of formula I



(I)


in which compound A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, for the preparation of a medicament for the treatment of a proliferative disease.

11. A commercial package comprising (a) an antidiarrheal agent and (b) an epothilone derivative of formula I



wherein A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, together with instructions for simultaneous, separate or sequential use thereof in the treatment of a proliferative disease.

12. A combined preparation, which comprises (a) one or more unit dosage forms of an antidiarrheal agent and (b) one or more unit dosage forms of an epothilone derivative of formula I



in which compound A represents O or NR_N , wherein R_N is hydrogen or lower alkyl, R is hydrogen or lower alkyl, and Z is O or a bond, in free form or in the form of a pharmaceutically acceptable salt.

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
26 September 2002 (26.09.2002)

PCT

(10) International Publication Number
WO 02/074042 A3

(51) International Patent Classification⁷: **A61K 31/445**,
33/24, 33/06, 38/31, 31/427, A61P 1/12, 35/00 // (A61K
31/427, 31:445, 33:06, 33:24, 38:31)

[US/US]; 46 Lakeview Terrace, Watchung, NJ 07069
(US). **TIANLING CHEN, TianLing** [US/US]; 38 Park
Street, #18C, Florham Park, NJ 07932 (US).

(21) International Application Number: PCT/EP02/02977

(74) Agent: **BECKER, Konrad**; Novartis AG, Corporate
Intellectual Property, Patent & Trademark Department,
CH-4002 Basel (CH).

(22) International Filing Date: 18 March 2002 (18.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/277,153 19 March 2001 (19.03.2001) US
60/277,207 20 March 2001 (20.03.2001) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LT, LU, LV, MA, MD, MK, MN, MX, NO, NZ, OM, PH,
PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TN, TR, TT, UA,
US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except AT, US*): **NO-
VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel
(CH).

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(71) Applicant (*for AT only*): **NOVARTIS-ERFINDUNGEN
VERWALTUNGSGESELLSCHAFT M.B.H.** [AT/AT];
Brunner Strasse 59, A-1230 Vienna (AT).

Published:
— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ROTHERMEL,
John, David** [US/US]; 46 School House Road, Randolph,
NJ 07869 (US). **SCHRAN, Horst, F.** [US/US]; 17 Junard
Drive, Morristown, NJ 07960 (US). **GREELEY, Diane**

(88) Date of publication of the international search report:
27 February 2003

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/074042 A3

(54) Title: COMBINATIONS COMPRISING AN ANTIDIARRHEAL AGENT AND AN EPOTHILONE OR AN EPOTHILONE
DERIVATIVE

(57) Abstract: Epothilone derivatives are co-administered with an antidiarrheal agent, e.g., a DPP-IV inhibitor, in the treatment of
a proliferative disease.

INTERNATIONAL SEARCH REPORT

 Interna application No
 PCT/EP 02/02977

A. CLASSIFICATION OF SUBJECT MATTER

 IPC 7 A61K31/445 A61K33/24 A61K33/06 A61K38/31 A61K31/427
 A61P1/12 A61P35/00 //(A61K31/427, 31:445, 33:06, 33:24, 38:31)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE, BIOSIS, EMBASE, PASCAL, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO 01 92255 A (KOSAN BIOSCIENCES INC ;ASHLEY GARY (US); FARDIS MARIA (US); SANTII) 6 December 2001 (2001-12-06) page 2, line 10 -page 3, line 9 page 40, line 19 - line 30	1-12
P,A	ALTAHA RAMIN ET AL: "Epothilones: A Novel Class of Non-taxane Microtubule-stabilizing Agents." CURRENT PHARMACEUTICAL DESIGN. NETHERLANDS 2002, vol. 8, no. 19, 2002, pages 1707-1712, XP001119806 ISSN: 1381-6128 Section "Clinical Experience", Pages 1709-1710	1-12

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

25 November 2002

Date of mailing of the international search report

05/12/2002

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Villa Riva, A

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 02/02977

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 5,6 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internal
Publication No
PCT/EP 02/02977

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0192255 A	06-12-2001	AU 6658301 A	11-12-2001
		WO 0192255 A2	06-12-2001
		US 2002045609 A1	18-04-2002
<hr/>			

THIS PAGE BLANK (USPTO)